

## DATA SHEET

<b>Catalog #</b>	AG-10200-228
<b>Cell Line Designation</b>	Parathyroid Hormone Receptor 2 cell line
<b>Parental Cell</b>	HEK 293-CNG cell (AG-10200-200)
<b>Gene Introduced</b>	Human Parathyroid Hormone Receptor 2 (PTHr2)
<b>NCBI Accession #</b>	NP_005039

### USAGE

- cAMP assay for Gs-coupled human Parathyroid Hormone Receptor 2 (PTHr2).
- HEK293-CNG cells (AG-10200-200) without transfected Parathyroid Hormone Receptor 2 are used as a negative control.


### QUALITY CONTROL

1. This cell line has been tested negative for *Mycoplasma sp.*
2. This cell line has been tested positive for Parathyroid Hormone Receptor 2 specific response.
3. Surviving rate: More than 2.5 million/vial on the second day after thawing.
4. The receptor specific activity is stable for 10 weeks continuous passage.

### CELL CULTURE CONDITION

1. Growth medium: 90% DMEM, 10% FBS, 250 µg/ml G418 and 1 µg/ml puromycin
2. Freezing medium: 10% DMSO, 90% complete medium

### DATA EXAMPLE



#### Figure 1. Response of ACTOne PTHR2 cell line & parental cell line to PTH.

ACTOne PTHR2 cells and parental cells (AG-10200-200) were plated overnight in 20 ml culture medium on a BD Biocoat 384 well plate. The next day, cells were dye-loaded with 20 ml/well of 1X Dye-loading solution (ACTOne Membrane Potential Assay Kit). After 2 hours of incubation at room temperature, two readings were obtained prior to and 30 min after the addition of PTH. Ratios of the two readings (F/F0) are plotted in the figure.

- A. Dose response curve of PTH in ACTOne PTHR2 cell line. EC50 = 20 pM in the presence of PDE inhibitor Ro20-1724, and EC50 = 73.7 pM in the absence of Ro20-1724 (data not shown).**
- B. Parental cells do not respond to PTH.**