

DATA SHEET

Catalog #	AG-10300-232
Cell Line Designation	Adenosine A3 receptor cell line
Parental Cell	HEK 293-CNG cell (AG-10200-200)
Gene Introduced	Human Adenosine A3 receptor (ADORA3)
NCBI Accession #	NP_000668

USAGE

- cAMP assay for Gi-coupled human Adenosine A3 receptor.
- HEK293-CNG cells (AG-10200-200) without transfected Adenosine A3 receptor are used as a negative control.

QUALITY CONTROL


1. This cell line has been tested negative for *Mycoplasma sp.*
2. This cell line has been tested positive for Adenosine A3 receptor specific response.
3. Surviving rate: More than 2.5 million/vial on the second day after thawing.
4. The receptor specific activity is stable for 10 weeks continuous passage.

CELL CULTURE CONDITION

1. Growth medium: 90% DMEM, 10% FBS, 250 $\mu\text{g/ml}$ G418 and 1 $\mu\text{g/ml}$ puromycin
2. Freezing medium: 10% DMSO, 90% complete medium

DATA EXAMPLE

A



B




Figure 1. Response of ACTOne Adenosine A3 receptor cell line & parental cell line to IB-Meca.

ACTOne Adenosine A3 receptor cells and parental cells (AG-10200-200) were plated overnight in 20 ml culture medium on a BD Biocoat 384 well plate. The next day, cells were dye-loaded with 20 ml/well of 1X Dye-loading solution (ACTOne Membrane Potential Assay Kit). After 2 hours of incubation at room temperature, two readings were obtained prior to and 30 min after the addition of IB-Meca. Ratios of the two readings (F/F0) are plotted in the figure.

- A. Dose response curve of IB-Meca in ACTOne Adenosine A3 receptor cell line. EC50 = 9 nM in the presence of PDE inhibitor Ro20-1724 and b-adrenoceptor agonist isoproterenol.**
- B. Parental cells do not respond to IB-Meca.**